

Central Pollution Control Board
(Ministry of Environment, Forest and Climate Change)
Govt. of India, Delhi

IMPACT OF LOCKDOWN (25th March to 15th April) ON AIR QUALITY

The nationwide Lockdown, in effect since the midnight of 24th March in view of COVID-19 pandemic, has resulted in significant improvement in air quality in the country, as revealed by data analysis and comparison of data for time before enforcement of restrictions. The Lockdown was announced after a 14-hour voluntary curfew called Janata curfew which was observed on 22 March, and subsequently CPCB published a report titled "IMPACT OF JANTA CURFEW & LOCKDOWN ON AIR QUALITY" dated 31.03.2020, describing the air quality trends in the country. This report is in continuation of the same. The major sectors contributing to air pollution are transport, industries, power plants, construction activities, biomass & refuse burning, road dust resuspension and residential activities. In addition, certain activities such as operation of DG sets, restaurant, landfill fires, etc. also contribute to air pollution. Under the nationwide lockdown, all transport services – road, air and rail were suspended with exceptions for essential services. Educational institutions, industrial establishments and hospitality services were also suspended. As a result, air quality improvement has been noted in many towns and cities across the nation.


Data generated from continuous ambient air quality monitoring (CAAQM) network has been analysed for Delhi with 38 stations and its neighbouring major NCR towns i.e. Faridabad, Gurugram, Noida and Ghaziabad with 4 stations each for the period from 16 March 2020 to 15 April 2020. Further, CAAQM data has also been analysed for a few major metropolitan cities i.e. Mumbai and Bengaluru with 10 stations each, Kolkata with 7 stations and a city in the Indo-Gangetic Plain, Patna with 6 stations for the period from 16 March 2020 to 15 April 2020 so as to obtain air quality trends, which have been studied in two phases: Pre-lockdown phase (16-21 March 2020) and Lockdown phase (25 March- 15 April 2020). AQI values as per CPCB bulletin have also been analysed along with satellite data (Aerosol Optical Depth) to observe the general trend of air quality improvement in the country.




EFFECT OF LOCKDOWN IN DELHI


During the lockdown period, as a result of combination of reduced vehicles on the road, functioning of only essential commercial units and prevailing weather conditions, significant reduction in PM_{2.5}, PM₁₀ and NO₂ levels were observed. Overall, 46% reduction in PM_{2.5} and 50% reduction in PM₁₀ was observed during the lockdown period. Similar level of reduction in PM₁₀ & PM_{2.5} primarily indicate reduction in combustion and industrial sources which are common to both fractions of Particulate matter. Since 81% of Delhi's NO_x comes from the transport sector (as per TERI Emission Inventory, 2018), restrictions on vehicular activity led to a 56% reduction in NO₂ levels and over 37% reduction in CO levels during the lockdown period, compared pre- lockdown period. Since there are restrictions in place on the transport sector and over

industrial operations, the two major sources of Benzene emissions, 47% reduction in Benzene levels has been observed. However, only 19% reduction was seen in SO₂ levels which may be due to the fact that over 70% of Delhi's SO₂ originates from power plants located around Delhi (as per TERI Emission Inventory, 2018) and power plants were operational during lockdown period. Other sources of SO₂ include restaurants and some industries, which might be operational during the lockdown period along with biomass/refuse burning in some areas in and around Delhi. However, most of these eateries and industries in Delhi have shifted from coal to other less polluting energy sources and thus, power plants appear to be the most likely source of Delhi's SO₂. 24 Hourly Average PM_{2.5} and PM₁₀ were within National Ambient Air Quality Standards (NAAQS) for 20 and 15 days respectively in 22 days of the lockdown period, while NO₂ levels were 75% less than their 24 hourly standard during the lockdown period. PM₁₀, PM_{2.5}, NO₂ and SO₂, 24 hourly average levels dropped as low as 39 µg/m³, 24 µg/m³, 15 µg/m³ and 10 µg/m³ respectively during the lockdown period.



A 27% increase in PM_{2.5} and a 65% increase in PM₁₀ compared to the first two weeks of the lockdown (25 March to 6 April) was observed in the second week of April (7 April to 15 April). However, PM_{2.5} and PM₁₀ levels were still lower by 39% and 35% respectively than pre-lockdown concentrations. This may primarily be attributed to change in meteorological conditions. Due to the onset of summers, temperature has started to increase with a minimum and maximum temperature of 12.6 °C and 27 °C on 16th March 2020 to 24 °C and 40°C on 15th April 2020, leading to dry and dusty conditions. Moreover, it was reported that a mild dust storm from western part of the country and the gulf regions hit Delhi on 14- 15th April, thus rapidly increasing the PM₁₀ levels in Delhi and NCR









The graphs below depict hourly concentration trend for $\text{PM}_{2.5}$ and PM_{10} , for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).





The hourly comparison of average concentration values shows a declining trend in levels of PM₁₀ and PM_{2.5} during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM₁₀ was 244 µg/m³ at 22:00 Hrs, which dropped to 127 µg/m³ during the lockdown period. Similarly, the lowest concentration during the pre-lockdown period was 118 µg/m³ at 17:00 Hrs, which dropped to 63 µg/m³ during the lockdown period. The drop in coarse particles may be attributed to restriction on construction activities, less road dust resuspension and to some extent curb on industrial activities. A similar decline was seen for PM_{2.5} with concentration value falling from a peak of 114 µg/m³ at 08:00 Hrs (during the pre-lockdown period) to a minimum value of 21 µg/m³ at 17:00 Hrs during the lockdown period. The absence of non-essential vehicles and combustion activities in industrial and commercial sites during the period is attributable to the decline.

The graphs below depict hourly concentration trend for NO₂ and CO for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).



Hourly NO₂ and CO values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO₂ during the pre-lockdown period was more than twice the peak value observed during the lockdown period. Similarly, peak hourly CO value also decreased by 55% during the lockdown period. The routine diurnal variation of NO₂ and CO is twin-crested with a larger crest during night hours. During the lockdown period, the night crest is much reduced, highlighting the absence of vehicular emissions.


The graphs below depict hourly concentration trend for SO₂ and Benzene for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).



Hourly Benzene levels during the lockdown period remained below the hourly values observed during the pre-lockdown period. Due to the reduced vehicular activity and restrictions on industrial operations, peak benzene levels reduced by 49% while its minimum value reduced by over 50%. However, hourly SO₂ values during the lockdown period were almost similar to the pre-lockdown values in the early

morning hours, i.e. when mixing height layer is less and ventilation is reduced. Moreover, the peak hourly SO₂ value only reduced by about 14% as Delhi's SO₂ largely comes from the power plants operating in its vicinity and that power plants were operational during the lockdown period.

It is important to mention here, the impact of meteorological factors was partially favorable, with maximum mixing depth of 4980m during the lockdown period higher than 3200 m recorded in the pre-lockdown period. Wind speed was higher during the lockdown period (4.7 m/s) as compared to the pre-lockdown period (3.9 m/s). However, with increase in temperature due to onset of summers, with high wind speed, there is an increased possibility of localized lifting of dust, thereby negatively affecting air quality. This is also depicted in the PM_{2.5}/PM₁₀ Ratio graph below. PM_{2.5}/PM₁₀ ratio started decreasing after 5th April and has been largely below 0.5 since then. This implies that the coarser particle (dust) is playing a dominant role in Delhi's Air Quality. The ratio fell drastically after 10th April and almost reached 0.3 on 15th April, primarily due to a mild dust storm from the Gulf regions (reported by IITM) hitting Delhi, thus significantly increasing the PM₁₀ concentration in Delhi.





As reported in source apportionment study conducted by TERI & ARAI, 2018, during summers, dust & construction activities (35%), transport sector (20%) and industry (20%), are major source of particulate matter in Delhi. As result of complete restrictions on non-essential vehicular movement and commercial activities, the emissions from construction activities and industries were stopped. The on-road vehicles were relatively sparse compared to normal days thus contribution from road dust resuspension & transport sector was much reduced. As evident from monitored data, the PM₁₀ emissions and PM_{2.5} emissions were reduced by up to 50% and 46% respectively.


Air Pollution Hotspots in Delhi

The data analysis of the 13 hotspots of Delhi reveals that Anand Vihar recorded 62%, 69% and 72% reduction in PM_{2.5}, PM₁₀ and NO₂ levels respectively during the lockdown period, as compared to the pre-lockdown period. However, data availability was low. Vivek Vihar, which is near to GT Road, a major traffic corridor, saw 60% reduction in NO₂ levels. Similarly, 48%, 61% and 68% reduction was observed in

Dwarka Sector-8. Further, Okhla recorded 72% reduction in NO₂ levels. It is to be noted that while Anand Vihar is a major transport hub and Okhla is major industrial suburb, Dwarka has substantial presence of residential cum institutional sites with substantial traffic movements, thus sharp decline in NO₂ levels further affirm that traffic and industrial operation restrictions were instrumental in improving air quality.

The box plots given below depict average PM_{2.5}, PM₁₀ and NO₂ concentrations in the 13 hotspots in the pre-lockdown period (BL) and during the lockdown period (DL).





The plots indicate distribution of values or the standard deviation for all hotspots has decreased considerably, suggesting the absence of major emission sources which contribute to variation in pollutant levels. Median values were seen to decrease for all hotspots for PM_{2.5} and PM₁₀ but increased for Jahangirpuri in the case of NO₂. This could be attributed to the presence of several inter-state goods carriers in the vicinity due to a major national highway located nearby.

EFFECT OF LOCKDOWN IN MAJOR NCR TOWNS

The air pollution reduction trend in NCR towns was relatively less pronounced compared to NCT of Delhi. Over 48% Reduction in PM₁₀ and PM_{2.5} levels were observed during lockdown period in all neighboring towns with sharp improvement in Faridabad with 55% reduction in PM_{2.5} levels and Gurugram with 54% reduction in PM₁₀ levels While significant reduction in NO₂ levels was observed in Noida (68%), Ghaziabad (60%) and Gurugram (40%), the same was not noted in Faridabad (17%), where NO₂ emissions were found higher during a few days in lockdown period, seemingly due to the gas-based power plants in and around Faridabad. Significant reduction in SO₂ levels was only seen in Faridabad (47%) and Ghaziabad (22%), while Gurugram (14%) and Noida (10%) recorded slight reductions during the lockdown period which may be attributed to their proximity to thermal power plants and some operational industries in the vicinity. Moreover, while Delhi's industries have largely switched over to gas-based and other less polluting energy systems, some industries in NCR might still be using coal and biomass, etc. Overall, average CO levels decreased in all major NCR towns with peak hourly values decreasing by 56% in Faridabad, 52% in Noida and 48% in Gurugram indicating reduced vehicular exhaust impact during lockdown. In terms of Benzene levels, Noida and Ghaziabad recorded an enormous reduction of 60% and 90% respectively during the lockdown period. However, Faridabad saw increase in Benzene levels. The operation of certain units or


processes (chemical/pharmaceutical/paints) utilizing benzene, and other solvents, etc. in Faridabad cannot be ruled out and may be responsible for the increase in Benzene levels. It is also important to mention that there are lesser number of real time air quality monitoring stations in NCR towns as compared to Delhi and the impact of localized sources on air quality data is always a possibility which may require further data for complete analysis. Moreover, in absence of complete data on scale of industrial operation in various categories except power plants and essential activities like food, bakery, dairy etc, it may be difficult to assess the impact of these sources on air quality at this point of time.




The trends for neighboring NCR towns are presented in detail in subsequent paras,




FARIDABAD




The impact of restrictions was visible in Faridabad. 19 days in the 22-day lockdown period witnessed 24 hourly PM_{2.5} levels within National Ambient Air Quality Standards (NAAQS). However, SO₂ and NO₂ values remained within National standards during pre-lockdown and lockdown period. While there has been a big drop in the peak hourly values of PM_{2.5} (44%), PM₁₀ (60%), SO₂ (57%) and CO (56%) during the lockdown period as compared to the pre-lockdown period, the minimum hourly value of NO₂ dropped only by 10% and hourly NO₂ values roughly remained the same throughout the day, in all probability due to localized combustion activities and operation of gas-based power plants in the vicinity. However, 57% reduction in peak hourly SO₂ levels with the minimum hourly SO₂ levels also decreasing by 29% during the lockdown period indicates that SO₂ emissions from coal-based power plants in NCR might be playing a more dominant role in Delhi than Faridabad


The data trends for Faridabad is as presented below,


















GURUGRAM




PM_{2.5}, NO₂ and SO₂ levels remained below National Ambient Air Quality Standards on all days during the lockdown, while PM₁₀ levels were above NAAQS on just 4 days in the 22-day lockdown period. Hourly data reveals a declining trend in pollutant levels from 07:00 Hrs onwards for major pollutants. Further, since dust & construction activities contribute 49% to PM_{2.5} and 52% to PM₁₀ in Gurugram (TERI Source Apportionment study, 2018), it is likely that road dust resuspension due to vehicle restrictions might have come down resulting in lower emissions with 42% and 52% reduction in peak hourly PM_{2.5} and PM₁₀ levels respectively. Peak hourly CO values reduced by almost 48% while peak hourly NO₂ and benzene levels reduced by 42% and 36% respectively during the lockdown period, indicating reduced vehicle movement. Peak SO₂ values fell by 20%, i.e. on a similar scale as Delhi.


The data trend for Gurugram is as presented below,






















GHAZIABAD

SO₂ and NO₂ levels remained below National Ambient Air Quality Standards on all days during the lockdown, while PM_{2.5} levels were above NAAQS in the second week of April i.e. for 7 days in the 22-day lockdown period. Although PM_{2.5} and PM₁₀ levels were higher during early morning hours and late-night hours, characteristically due to reduced ventilation and mixing height, peak hourly PM_{2.5} and PM₁₀ levels reduced by 49% and 57% respectively. Major reduction of 91% in peak benzene levels and 66% in peak NO₂ values was observed, largely due to the reduced presence of vehicular and industrial activity. Major reduction in benzene levels (highest in Delhi NCR) during lockdown period indicate closure of some large-scale benzene utilizing/generating source like paint, petro products, plastics, resins, synthetic fibers, rubber lubricants, dyes, detergents, drugs and pesticides in Ghaziabad region apart from reduced impact of vehicular related emissions. While average CO values during lockdown period remained below their pre-lockdown levels to a great extent and reduced by almost 30%, analysis of hourly CO values indicate peak hourly CO value rising by 11%, seemingly due to local combustion activities which may include increased use of solid fuels/biomass in household cooking etc. Further, the diurnal cycle of CO concentration presents two peaks, in the morning and in the evening. Notably, mixing height is generally low in these two periods of the day. Further, studies indicate that in the early hours of the morning, surface heating by solar radiation is also not enough to break the previous night's thermal inversion layer, causing the pollutants to remain concentrated in regions close to the surface. As mixing height increases allowing transport of pollutants to the upper layers, CO levels on the surface decrease and rise again by the end of the afternoon, when convective activity decreases and traffic generally increases.

















NOIDA




Positive effects of lockdown on air pollution levels were observed in Noida, as emission levels considerably reduced from the pre-lockdown period with over 48% reduction in PM_{2.5} and PM₁₀. 24 hourly average PM_{2.5} and PM₁₀ concentrations remained within NAAQS for 19 and 17 days respectively out of the 22 days in the lockdown period. NO₂ and SO₂ levels remained within NAAQS on all days of the lockdown period with the peak hourly NO₂ value decreasing from 76 µg/m³ in the pre-lockdown period to 20 µg/m³ in the lockdown period. Peak hourly Benzene levels reduced by 67% , in all possibility due to the restrictions on vehicular activity and industrial operations. While construction activity is a major emission source in Noida contributing 47% to PM₁₀ (TERI source apportionment study, 2018) , significant reduction in PM₁₀ levels with hourly peak values decreasing by 55%, suggest reduced contribution of road dust resuspension & C & D activities. Reduction in PM_{2.5} and CO emission levels was lower during morning hours signifying contribution of combustion activities. Further, over 52% reduction in peak hourly CO values was observed.

While overall SO₂ levels were seen to decline during the lockdown period, peak hourly SO₂ value increased marginally. Hourly SO₂ levels were also higher during the early morning hours, when dispersion of pollutants is lower. It may be said that thermal power plants located in NCR and use of fuels like coal and biomass/wood etc in industrial and household activities including operation of some brick kilns, Sugar and distilleries, might be playing a more dominant role in affecting SO₂ levels in Noida.









EFFECT OF LOCKDOWN IN DELHI NCR

Substantial improvement in air quality of Delhi NCR is noted during the lockdown period, as the major contributing sources to PM & NO₂ emissions (prominent pollutants in Delhi NCR) have been restricted. The AQI in Delhi NCR was largely under 'moderate' category in the week before start of lockdown period. As days progressed, under cumulated effect of restricted vehicle movement, industrial & commercial activities and increased mixing height, the AQI improved to 'Satisfactory' category. On March 26, 2020, high surface winds (25 kmph) maintained AQI category even though mixing height dropped to 1100 m. Next day, though wind speed and mixing height were reduced to half value, AQI value improved further and Gurugram recorded 'Good' AQI category. Scattered rains in Delhi NCR on 27th March and during March 28- 29, 2020 along with increased wind speed and mixing height, AQI value improved further, with Delhi, Ghaziabad and Noida recording 'Good' AQI category on March 28, 2020. Favorable conditions ensued, leading to AQI remaining in 'Good' and 'Satisfactory' AQI categories. However, after 4th April due to change in temperature and onset of dry conditions, high winds led to lifting of local dust resulting in slight deterioration of air quality to 'moderate' category. Further, a dust storm from the gulf hit Delhi and the surrounding areas on 15th April, further pushing the air quality to the higher end of moderate category.



Date	Predominant Wind Speed (kmph)	Maximum Mixing Height (m)	Delhi	Ghaziabad	Noida	Faridabad	Gurugram
16-Mar	16	1800	139	134	118	184	165
17-Mar	12	1500	157	148	140	164	141
18-Mar	15	2000	151	172	137	164	168
19-Mar	14	3200	186	236	184	194	192
20-Mar	12	2400	192	235	195	212	175
21-Mar	16	2500	186	207	161	174	126
22-Mar (Janata Curfew)	12	2900	191	237	176	214	191
23-Mar	10	800	124	159	123	130	91
24-Mar	10	2700	122	166	130	187	127
IMPOSITION OF NATIONWIDE LOCKDOWN DUE TO COVID-19							
25-Mar	12	2500	77	86	80	100	69
26-Mar	25	1100	92	84	72	88	61
27-Mar	15	500	69	72	60	75	42
28-Mar	14	2250	45	39	38	64	54
29-Mar	20	2600	62	48	58	83	62
30-Mar	20	2100	71	64	61	97	76
31-Mar	12	1900	76	72	67	110	77
01-Apr	12	3200	73	79	73	90	69
02-Apr	20	3050	69	63	62	63	72
03-Apr	22	2100	79	104	72	97	82




04-Apr	20	3000	87	109	70	90	89
05-Apr	10	3500	102	124	84	117	91
06-Apr	15	3500	142	181	120	123	106
07-Apr	32	3750	90	101	78	100	86
08-Apr	20	3100	83	113	85	103	91
09-Apr	16	2400	86	86	80	103	96
10-Apr	15	3500	118	115	93	117	104
11-Apr	8	3050	124	194	146	203	152
12-Apr	20	2750	94	93	87	119	98
13-Apr	18	3700	126	132	120	122	106
14-Apr	12	4480	130	145	123	146	113
15-Apr	12	4980	155	194	184	186	142

AIR QUALITY TREND ANALYSIS IN OTHER CITIES DURING LOCKDOWN PERIOD



MUMBAI

During the lockdown period, significant reduction in PM_{2.5}, PM₁₀ and NO₂ levels were observed. Overall, 40% reduction in PM_{2.5} and 43% reduction in PM₁₀ was observed during the lockdown period. A sharp decrease of 77% reduction in NO₂ levels and 59% decrease in average Benzene levels was observed during the lockdown period, compared to the week before lockdown came into force, largely due to the reduced presence of vehicular activity and restricted industrial operations. Average SO₂ levels decreased by 33% in the lockdown period, largely due to shutdown of industries. As per emission inventory of Mumbai (CPCB, 2010), 39 types of industries (excluding power plants) contribute to over 50% of SO₂. 24 hourly average PM_{2.5}, PM₁₀, SO₂ and NO₂ levels were within National Ambient Air Quality Standards for all days in the lockdown period. 24 hourly average PM₁₀, PM_{2.5} and NO₂ levels dropped as low as 54 µg/m³, 20 µg/m³ and 5 µg/m³ respectively during the lockdown period.







The graphs below depict hourly concentration trend for PM_{2.5} and PM₁₀, for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).



The hourly comparison of average concentration values shows a declining trend in levels of PM₁₀ and PM_{2.5} during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM₁₀ was 175 µg/m³ at 08:00 Hrs, which dropped to 85 µg/m³ during the lockdown period. Similarly, the lowest concentration during the pre-lockdown period at 17:00 Hrs was 98 µg/m³, which dropped to 58 µg/m³ during the lockdown period, seemingly due to restriction on construction activities, less road dust resuspension and to some extent curb on industrial activities. A similar decline was seen for PM_{2.5} with concentration value falling from a peak of 62 µg/m³ at 08:00 Hrs (during the pre-lockdown period) to a minimum value of 21 µg/m³ at 05:00 Hrs during the lockdown period. The absence of non-essential vehicles and combustion activities in industrial and commercial sites during the period may be attributable to the decline. Notably, the



lowest values are occurring in early morning hours during the lockdown period against a deeper crest in evening hours during the pre-lockdown period. This indicates the absence of nighttime accumulation of pollutants, highlighting the absence of major emission sources.

The graphs below depict hourly concentration trend for NO₂ and CO for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).



Hourly NO₂ and CO values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO₂ during the pre-lockdown period was almost five times the peak value observed during the lockdown period. Similar trend was observed for CO with 61% reduction in peak hourly values during the lockdown period, underscoring the absence of vehicular emissions.




The graphs below depict hourly concentration trend for SO₂ and Benzene for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).





As a result of the lockdown, hourly Benzene and SO₂ levels remained below their pre-lockdown levels at all times. The diurnal variation in the pollutants was reduced, implying the reduced impact of major emission sources. Peak hourly SO₂ values declined by 46% while peak hourly Benzene Concentration reduced by over 70%, seemingly due to absence of vehicular activity.


KOLKATA


During the lockdown period, over 40% reduction in PM_{2.5}, 44% in PM₁₀ levels and 38% reduction in SO₂ levels were observed. CO average levels also decreased in a similar manner with 39% reduction during the lockdown period. However, 66% reduction in NO₂ levels and 69% reduction in Benzene levels was observed during the lockdown period, indicating the reduction in number of on-road vehicles. 24 hourly average PM_{2.5}, PM₁₀, SO₂ and NO₂ levels were within National Ambient Air Quality Standards for all days in the lockdown period. PM₁₀, PM_{2.5} and NO₂ 24 hourly average levels dropped as low as 38 µg/m³, 18 µg/m³ and 6 µg/m³ during the lockdown period.






The graphs below depict hourly concentration trend for PM_{2.5} and PM₁₀, for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).






The hourly comparison of average concentration values shows a declining trend in levels of PM₁₀ and PM_{2.5} during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM₁₀ was 150 µg/m³ at 00:00 Hrs, which dropped to 73 µg/m³ during the lockdown period. Similarly, the lowest concentration during the pre-lockdown period at 16:00 Hrs was 92 µg/m³, which dropped to 56 µg/m³ during the lockdown period. The drop in coarse particles may be attributed to restriction on construction activities, less road dust resuspension and to some extent curb on industrial activities. A similar decline was seen for PM_{2.5} with concentration value falling from a peak of 80 µg/m³ at 00:00 Hrs (during the pre-lockdown period) to a minimum value of 33 µg/m³ at 11:00 Hrs during the lockdown period which may be attributed to the absence of non-essential vehicles and combustion activities in industrial and commercial sites. Kolkata's diurnal variation of pollutants is characterized by lower concentrations during the day and accumulation of pollutants during nighttime which was reduced due to the absence of daytime activities like vehicular movement and industrial operation.


The graphs below depict hourly concentration trend for NO₂ and CO for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).






Hourly NO₂ and CO values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO₂ during the pre-lockdown period was more than thrice the peak value observed during the lockdown period. Similar trend was observed for CO, with hourly peak CO values reducing by almost 59%.

The graphs below depict hourly concentration trend for SO₂ and Benzene for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).










Due to restrictions on industrial operations and vehicular activity, SO_2 and Benzene levels remained below their pre-lockdown levels during the lockdown period. A decrease of 49% was observed in peak hourly SO_2 values. There was little diurnal variation in Benzene levels with over 77% reduction in peak hourly Benzene values while hourly SO_2 levels increased as the day progressed, which may be attributed to the thermal power plants running in and around Kolkata along with local combustion activities.



BENGALURU

Significant reduction in PM_{10} and NO_2 levels were observed during the lockdown period, as a result of combination of reduced vehicles on the road and functioning of only essential commercial units. Overall, 44% reduction in PM_{10} and 21% reduction in $\text{PM}_{2.5}$ was observed during the lockdown period. Higher reduction in PM_{10} levels indicate closure of dust generating activities like construction and Demolition etc. Due to the restrictions imposed on vehicular movement and industrial activity, a 53% reduction in NO_2 levels and 48% reduction in Benzene levels was observed during the lockdown period, compared to the week before lockdown came into force. 24 hourly average $\text{PM}_{2.5}$, PM_{10} , SO_2 and NO_2 levels were within National Ambient Air Quality Standards for all days in the lockdown period. 24 hourly average PM_{10} , $\text{PM}_{2.5}$ and NO_2 levels dropped as low as $20 \mu\text{g}/\text{m}^3$, $15 \mu\text{g}/\text{m}^3$ and $9 \mu\text{g}/\text{m}^3$ during the lockdown period. However, only 15% reduction was seen in SO_2 levels during the lockdown period. It is possible that use of fuels like coal and biomass/wood etc in industrial and household activities, might be influencing SO_2 levels in Bengaluru. Similar reduction was seen in CO levels too, seemingly due to local combustion activities.







The graphs below depict hourly concentration trend for PM_{2.5} and PM₁₀, for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).





The hourly comparison of average concentration values shows a declining trend in levels of PM₁₀ and PM_{2.5} during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM₁₀ was 135 µg/m³ at 20:00 Hrs, which dropped to 56 µg/m³ during the lockdown period. Similarly, the lowest hourly concentration during the pre-lockdown period at 03:00 Hrs was 64 µg/m³, which dropped to 37 µg/m³ during the lockdown period, in all likelihood due to restriction on industrial operations, construction activities and reduced road dust resuspension. A decline was also seen for PM_{2.5} with concentration value falling from a peak of 46 µg/m³ at 20:00 Hrs (during the pre-lockdown period) to a minimum value of 18 µg/m³ at 14:00 Hrs during the lockdown period, possibly due to the reduced presence of vehicles and industrial/commercial combustion activities.

The graphs below depict hourly concentration trend for NO₂ and CO for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).



Hourly NO₂ and CO values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO₂ during the pre-lockdown period was thrice the peak value observed during the lockdown period. Diurnal variation in NO₂ has reduced due to decreased vehicular activity. Hourly CO values also remained below their pre-lockdown values for most of the hours except during early morning, with hourly peak value decreasing by 30%. Post lockdown, higher CO values during night hours have reduced and maximum concentration of 1.4 mg/m³ was observed at 20.00 hours before lockdown and 0.99 mg/m³ at 09.00 hours during lockdown.

The graphs below depict hourly concentration trend for SO₂ and Benzene for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).








Peak hourly Benzene levels recorded a decrease of 47% in the lockdown period, largely due to the reduction in the number of on-road vehicles. However, there was only a 9% reduction in peak SO₂ values. Hourly SO₂ values during the lockdown period appear to follow a similar pattern as pre-lockdown levels, implying that the major sources of SO₂ in Bengaluru might still be operational, despite the lockdown and requires further investigation.

PATNA



Overall, 27% reduction in PM_{2.5}, 27% reduction in PM₁₀ and 30% reduction in NO₂ levels was observed during the lockdown period, compared to the week before lockdown came into force. SO₂ and NO₂ levels remained within National Ambient Air Quality Standards for all of the days in the lockdown period. While PM_{2.5} levels were over NAAQS on just three days, PM₁₀ levels remained within NAAQS for only 5 days in the 22-day lockdown period . Since major source of PM₁₀ for Patna is dust, it is difficult to assess its

contribution during the period without additional data like meteorology. PM_{10} , $PM_{2.5}$ and NO_2 24 hourly average levels dropped to $84 \mu\text{g}/\text{m}^3$, $28 \mu\text{g}/\text{m}^3$ and $17 \mu\text{g}/\text{m}^3$ during the lockdown period. Benzene levels dropped by almost 61% in the lockdown period, indicating the absence of major industrial activities utilizing/emitting benzene. Although lower pollutant concentrations were observed in Patna, major reduction was not seen probably due to continuing movement of vehicles. CO and SO_2 levels, however were observed to slightly increase (5%), seemingly due to increased local combustion activities and usage of solid fuels in household and industrial units. However more information may be required about Sulphur dioxide sources in and around the city to examine the reasons behind the trend of SO_2 during lockdown.







The graphs below depict hourly concentration trend for PM_{2.5} and PM₁₀, for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).





The hourly comparison of average concentration values shows a declining trend in levels of PM₁₀ and PM_{2.5} during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM₁₀ was 220 µg/m³ at 22:00 Hrs, which dropped to 159 µg/m³ during the lockdown period. Similarly, the lowest concentration during the pre-lockdown period at 15:00 Hrs was 117 µg/m³, which slightly reduced to 92 µg/m³ during the lockdown period. A similar decline was seen for PM_{2.5} with concentration value falling from a peak of 84 µg/m³ at 23:00 Hrs (during the pre-lockdown period) to a minimum value of 32 µg/m³ at 17:00 Hrs during the lockdown period, probably due to the absence of non-essential vehicles and industrial operations during the period.

The graphs below depict hourly concentration trend for NO₂ and CO for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).



Hourly NO₂ and CO values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO₂ during the pre-lockdown period was over one and half times the peak value observed during the lockdown period. Though 11% reduction in hourly peak CO values was observed in the lockdown period, overall levels of CO were seen to increase, probably due to local combustion activities like biomass burning and agri-waste burning in surrounding areas.

The graphs below depict hourly concentration trend for SO₂ and Benzene for pre-lockdown period (16th March 2020 to 21st March 2020) and lockdown period (25th March 2020 to 15th April 2020).



While peak hourly Benzene value was seen to decrease by 61% in the lockdown period, peak hourly SO₂ value increased by 3%. Hourly SO₂ values during the lockdown period appear to be following the same diurnal variation as in the pre-lockdown period, implying that the major sources of SO₂, i.e. biomass burning or usage of unclean fuels in household and industrial activities including brick kilns might still be operational. The trend of SO₂ need to be further investigated with more information on likely sources including on-field data, as it is defying general trend observed during lockdown.

EFFECT OF LOCKDOWN IN OTHER CITIES IN TERMS OF AIR QUALITY INDICES

Air Quality Indices are calculated for cities all over India using data from CAAQM stations. If cities appearing in CPCB AQI Bulletin are grouped according to their respective AQI categories, it is observed that about 78% of cities in the AQI bulletin are falling in Good and Satisfactory categories in the lockdown phase, increasing from the average of 44% seen in the pre-lockdown phase. Since Good and Satisfactory categories have their breakpoints within the National Ambient Air Quality Standards, it may be reasonable to state that more cities have their air quality within National standards during the lockdown period.

During the lockdown period, no city entered the very poor category. Among the cities in poor category during the lockdown period, instances of Singrauli and Brajrajnagar are found frequently. It is worth noting that Singrauli region is home to several power plants, which are operational during the lockdown period and Brajrajnagar has in its vicinity numerous open-cast and underground coal mines.

The date wise AQI is given in Annexure I.

IMPACT OF LOCKDOWN (18th March to 13th April) ON AIR QUALITY – A SATELLITE BASED APPROACH

Satellite-based monitoring is being used widely for air quality data generation. The advantage of satellite-based estimates is that they can provide country wide coverage of all locations where people live in a consistent way.

A project entitled “Satellite based near real time monitoring of ambient PM_{2.5} at national scale for air quality management” has been initiated by CPCB in collaboration with Indian Institute of Technology (IIT), Delhi under National Clean Air Programme (NCAP) in order to strengthen development of indigenous satellite-based products and techniques to derive useful air quality information and to supplement the current monitoring network.

Satellites retrieve a parameter called aerosol optical depth (AOD), which represents columnar concentration of particulate matter in terms of light extinction. AOD is then converted to surface PM_{2.5} (fine particulate matter smaller than 2.5 µm) using a dynamic scaling factor that is modulated by vertical distribution of aerosols in the atmosphere, meteorological conditions and emission pattern. An algorithm of estimating PM_{2.5} from AOD product of Multi-Angle Implementation of Atmospheric Correction (MAIAC) onboard TERRA and Aqua satellites has been developed to map surface PM_{2.5} over India at 1-km resolution from 2000 onwards. The satellite-based 24-hr and annual PM_{2.5} show correlation of 0.87 and 0.98 and RMSE of 25 µg/m³ and 1.9 µg/m³ respectively with coincident in-situ data from CPCB network.

Impact of lockdown (Starting on March 25, 2020) on PM_{2.5} levels at a national scale was carried out using this unique data for the period March 18 to April 13, 2020. Weekly PM_{2.5} concentration anomaly (in %) in 2020 has been reported relative to 2015-2019 average concentration (see Figure). The analysis suggests that PM_{2.5} was higher in North and Central India during the pre-lockdown week this year. With many sources shut down, in the first two weeks of lockdown, PM_{2.5} was 30-50% lower in most parts of India, especially in the Indo-Gangetic Plain (IGP). The reverse signal in the Central India during 25th – 31st March may be attributed to intense open burning, which subsided in the week after as demonstrated by a negative anomaly. Similar trends are reported by few recent studies using CPCB in-situ data.

However, this improvement seems to be gone in the 2nd week of April, when the dust influence started to dominate. Data after 13th April is being analysed to understand whether this is temporary trend, however, ground-based data also showed that air quality is again back in moderate zone.




Figure. Anomaly (in % w.r.t last 5-years) in weekly surface PM_{2.5} from satellite-derived data during pre- and lockdown period

SUMMARY

During the lockdown period, there has been a general improvement in air quality in the country as a result of the restrictions imposed during the lockdown, which is corroborated by both Air Quality Index data and Satellite data. All of the major cities, analysed in this report, had their AQI, largely within Good-Moderate categories, with Bengaluru and Mumbai recording all 22 days of the lockdown period in Good-Satisfactory categories while Patna had majority of days in moderate category. This is of importance because most of these cities apart from being densely populated, have also been declared non-attainment cities (cities exceeding National Ambient Air Quality Standards).

**Comparative AQI Status from 16 March to 15 April, 2020
(based on CPCB AQI Bulletin, published at 4 PM)**

Date	No of cities for which data is available	No. of cities in AQI category						No. of cities with AQI in range of Good to Satisfactory	No. Of cities with AQI in Moderate Category	No. of cities with AQI in range of Poor to Severe
		Good	Satisfactory	Moderate	Poor	Very Poor	Severe			
16-Mar-20	108	6	49	50	3	0	0	55	50	3
17-Mar-20	111	3	44	59	5	0	0	47	59	5
18-Mar-20	112	3	42	58	9	0	0	45	58	9
19-Mar-20	115	3	39	65	8	0	0	42	65	8
20-Mar-20	115	2	51	50	12	0	0	53	50	12
21-Mar-20	112	2	52	49	9	0	0	54	49	9
22-Mar-20 (Janata Curfew)	114	9	58	39	8	0	0	67	39	8
23-Mar-20	108	10	63	33	2	0	0	73	33	2
24-Mar-20	110	11	54	43	2	0	0	65	43	2
National Lockdown in effect due to COVID-19 Pandemic										
25-Mar-20	104	14	67	21	2	0	0	81	21	2
26-Mar-20	102	21	64	14	3	0	0	85	14	3
27-Mar-20	103	31	59	10	3	0	0	90	10	3
28-Mar-20	101	35	57	8	1	0	0	92	8	1
29-Mar-20	103	30	61	12	0	0	0	91	12	0
30-Mar-20	99	23	65	11	0	0	0	88	11	0
31-Mar-20	103	20	67	14	2	0	0	87	14	2
01-Apr-20	100	23	62	13	2	0	0	85	13	2
02-Apr-20	105	22	71	11	1	0	0	93	11	1

03-Apr-20	105	20	71	14	0	0	0	91	14	0
04-Apr-20	109	22	68	18	1	0	0	90	18	1
05-Apr-20	104	17	65	21	1	0	0	82	21	1
06-Apr-20	102	23	49	29	1	0	0	72	29	1
07-Apr-20	101	25	56	18	2	0	0	81	18	2
08-Apr-20	102	22	54	25	1	0	0	76	25	1
09-Apr-20	102	20	58	24	0	0	0	78	24	0
10-Apr-20	104	17	54	31	2	0	0	71	31	2
11-Apr-20	103	21	49	29	4	0	0	70	29	4
12-Apr-20	108	17	62	28	1	0	0	79	28	1
13-Apr-20	104	15	50	38	1	0	0	65	38	1
14-Apr-20	102	8	53	36	5	0	0	61	36	5
15-Apr-20	105	8	49	38	10	0	0	57	38	10

AQI Category	AQI Range	Associated Health Impact
Good	0-50	Minimal Impact
Satisfactory	51-100	Minor breathing discomfort to sensitive people
Moderate	101-200	Breathing discomfort to the people with lungs, asthma and heart diseases
Poor	201-300	Breathing discomfort to most people on prolonged exposure
Very Poor	301-400	Respiratory illness on prolonged exposure
Severe	401-500	Affects healthy people and seriously impacts those with existing diseases

Number of cities with AQI in Good, Satisfactory and Moderate categories

