

**Central Pollution Control Board**  
**(Ministry of Environment, Forest and Climate Change)**  
**Govt. of India, Delhi**

## **IMPACT OF LOCKDOWN (25<sup>th</sup> March to 15<sup>th</sup> April) ON AIR QUALITY**

The nationwide Lockdown, in effect since the midnight of 24<sup>th</sup> March in view of COVID-19 pandemic, has resulted in significant improvement in air quality in the country, as revealed by data analysis and comparison of data for time before enforcement of restrictions. The Lockdown was announced after a 14-hour voluntary curfew called Janata curfew which was observed on 22 March, and subsequently CPCB published a report titled "IMPACT OF JANTA CURFEW & LOCKDOWN ON AIR QUALITY" dated 31.03.2020, describing the air quality trends in the country. This report is in continuation of the same. The major sectors contributing to air pollution are transport, industries, power plants, construction activities, biomass & refuse burning, road dust resuspension and residential activities. In addition, certain activities such as operation of DG sets, restaurant, landfill fires, etc. also contribute to air pollution. Under the nationwide lockdown, all transport services – road, air and rail were suspended with exceptions for essential services. Educational institutions, industrial establishments and hospitality services were also suspended. As a result, air quality improvement has been noted in many towns and cities across the nation.


Data generated from continuous ambient air quality monitoring (CAAQM) network has been analysed for Delhi with thirty-eight (38) stations and its neighbouring major NCR towns i.e. Faridabad, Gurugram, Noida and Ghaziabad with four (4) stations each for the period from 16 March 2020 to 15 April 2020. Further, CAAQM data has also been analysed for a few major metropolitan cities i.e. Mumbai and Bengaluru with ten (10) stations each, Kolkata with seven (7) stations, Chennai with four (4) stations, Indore with one (1) station and a city in the Indo-Gangetic Plain, Patna with 6 stations for the period from 16 March 2020 to 15 April 2020 so as to obtain air quality trends, which have been studied in two phases: Pre-lockdown phase (16-21 March 2020) and Lockdown phase (25 March- 15 April 2020). AQI values as per CPCB bulletin have also been analysed along with satellite data to observe the general trend of air quality improvement in the country.




### **EFFECT OF LOCKDOWN IN DELHI**


During the lockdown period, as a result of combination of reduced vehicles on the road, functioning of only essential commercial units and prevailing weather conditions, significant reduction in PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub> levels were observed. Overall, 46% reduction in PM<sub>2.5</sub> and 50% reduction in PM<sub>10</sub> was observed during the lockdown period. Similar level of reduction in PM<sub>10</sub> & PM<sub>2.5</sub> primarily indicate reduction in combustion and industrial sources which are common to both fractions of Particulate matter. Since 81% of Delhi's NO<sub>x</sub> comes from the transport sector (as per TERI Emission Inventory, 2018), restrictions on vehicular activity led to a 56% reduction in NO<sub>2</sub> levels and over 37% reduction in CO levels during the lockdown period,

compared pre- lockdown period. Since there are restrictions in place on the transport sector and over industrial operations, the two major sources of Benzene emissions, 47% reduction in Benzene levels has been observed. However, only 19% reduction was seen in SO<sub>2</sub> levels which may be due to the fact that over 70% of Delhi's SO<sub>2</sub> originates from power plants located around Delhi (as per TERI Emission Inventory, 2018) and power plants were operational during lockdown period. Other sources of SO<sub>2</sub> include restaurants and some industries, which might be operational during the lockdown period along with biomass/refuse burning in some areas in and around Delhi. However, most of these eateries and industries in Delhi have shifted from coal to other less polluting energy sources and thus, power plants appear to be the most likely source of Delhi's SO<sub>2</sub>. 24 Hourly Average PM<sub>2.5</sub> and PM<sub>10</sub> were within National Ambient Air Quality Standards (NAAQS) for 20 and 15 days respectively in 22 days of the lockdown period, while NO<sub>2</sub> levels were 75% less than their 24 hourly standard during the lockdown period. Lowest PM<sub>10</sub>, PM<sub>2.5</sub>, NO<sub>2</sub> and SO<sub>2</sub>, 24 hourly average levels during the lockdown period were recorded as 39 µg/m<sup>3</sup>, 24 µg/m<sup>3</sup>, 15 µg/m<sup>3</sup> and 10 µg/m<sup>3</sup> respectively.



A 27% increase in PM<sub>2.5</sub> and a 65% increase in PM<sub>10</sub> compared to the first two weeks of the lockdown (25 March to 6 April) was observed in the second week of April (7 April to 15 April). However, PM<sub>2.5</sub> and PM<sub>10</sub> levels were still lower by 39% and 35% respectively than pre-lockdown concentrations. This may primarily be attributed to change in meteorological conditions. Due to the onset of summers, temperature has started to increase with a minimum and maximum temperature of 12.6 °C and 27 °C on 16<sup>th</sup> March 2020 to 24 °C and 40°C on 15<sup>th</sup> April 2020, leading to dry and dusty conditions. Moreover, it was reported that a mild dust storm from western part of the country and the gulf regions hit Delhi on 14- 15<sup>th</sup> April, thus rapidly increasing the PM<sub>10</sub> levels in Delhi and NCR









The graphs below depict hourly concentration trend for  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$ , for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).





The hourly comparison of average concentration values shows a declining trend in levels of PM<sub>10</sub> and PM<sub>2.5</sub> during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM<sub>10</sub> was 244 µg/m<sup>3</sup> at 22:00 Hrs, which dropped to 127 µg/m<sup>3</sup> during the lockdown period. Similarly, the lowest concentration during the pre-lockdown period was 118 µg/m<sup>3</sup> at 17:00 Hrs, which dropped to 63 µg/m<sup>3</sup> during the lockdown period. The drop in coarse particles may be attributed to restriction on construction activities, less road dust resuspension and to some extent curb on industrial activities. A similar decline was seen for PM<sub>2.5</sub> with concentration value falling from a peak of 114 µg/m<sup>3</sup> at 08:00 Hrs (during the pre-lockdown period) to a minimum value of 21 µg/m<sup>3</sup> at 17:00 Hrs during the lockdown period. The absence of non-essential vehicles and combustion activities in industrial and commercial sites during the period is attributable to the decline.

The graphs below depict hourly concentration trend for NO<sub>2</sub> and CO for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).



Hourly NO<sub>2</sub> and CO values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO<sub>2</sub> during the pre-lockdown period was more than twice the peak value observed during the lockdown period. Similarly, peak hourly CO value also decreased by 55% during the lockdown period. The routine diurnal variation of NO<sub>2</sub> and CO is twin-crested with a larger crest during night hours. During the lockdown period, the night crest is much reduced, highlighting the absence of vehicular emissions.


The graphs below depict hourly concentration trend for SO<sub>2</sub> and Benzene for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).



Hourly Benzene levels during the lockdown period remained below the hourly values observed during the pre-lockdown period. Due to the reduced vehicular activity and restrictions on industrial operations, peak benzene levels reduced by 49% while its minimum value reduced by over 50%. However, hourly SO<sub>2</sub> values during the lockdown period were almost similar to the pre-lockdown values in the early

morning hours, i.e. when mixing height layer is less and ventilation is reduced. Moreover, the peak hourly SO<sub>2</sub> value only reduced by about 14% as Delhi's SO<sub>2</sub> largely comes from the power plants operating in its vicinity and that power plants were operational during the lockdown period.

It is important to mention here, the impact of meteorological factors was partially favorable, with maximum mixing depth of 4980m during the lockdown period higher than 3200 m recorded in the pre-lockdown period. Wind speed was higher during the lockdown period (4.7 m/s) as compared to the pre-lockdown period (3.9 m/s). However, with increase in temperature due to onset of summers, with high wind speed, there is an increased possibility of localized lifting of dust, thereby negatively affecting air quality. This is also depicted in the PM<sub>2.5</sub>/PM<sub>10</sub> Ratio graph below. PM<sub>2.5</sub>/PM<sub>10</sub> ratio started decreasing after 5<sup>th</sup> April and has been largely below 0.5 since then. This implies that the coarser particle (dust) is playing a dominant role in Delhi's Air Quality. The ratio fell drastically after 10<sup>th</sup> April and almost reached 0.3 on 15<sup>th</sup> April, primarily due to a mild dust storm from the Gulf regions (reported by IITM) hitting Delhi, thus significantly increasing the PM<sub>10</sub> concentration in Delhi.





As reported in source apportionment study conducted by TERI & ARAI, 2018, during summers, dust & construction activities (35%), transport sector (20%) and industry (20%), are major source of particulate matter in Delhi. As result of complete restrictions on non-essential vehicular movement and commercial activities, the emissions from construction activities and industries were stopped. The on-road vehicles were relatively sparse compared to normal days thus contribution from road dust resuspension & transport sector was much reduced. As evident from monitored data, the PM<sub>10</sub> emissions and PM<sub>2.5</sub> emissions were reduced by up to 50% and 46% respectively.

#### Air Pollution Hotspots in Delhi


The data analysis of the 13 hotspots of Delhi reveals that Anand Vihar recorded 62%, 69% and 72% reduction in PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub> levels respectively during the lockdown period, as compared to the pre-lockdown period. However, data availability was low. Vivek Vihar, which is near to GT Road, a major traffic corridor, saw 60% reduction in NO<sub>2</sub> levels. Similarly, 48%, 61% and 68% reduction was observed in

Dwarka Sector-8. Further, Okhla recorded 72% reduction in NO<sub>2</sub> levels. It is to be noted that while Anand Vihar is a major transport hub and Okhla is major industrial suburb, Dwarka has substantial presence of residential cum institutional sites with substantial traffic movements, thus sharp decline in NO<sub>2</sub> levels further affirm that traffic and industrial operation restrictions were instrumental in improving air quality.

The box plots given below depict average PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub> concentrations in the 13 hotspots in the pre-lockdown period (BL) and during the lockdown period (DL).







The plots indicate distribution of values or the standard deviation for all hotspots has decreased considerably, suggesting the absence of major emission sources which contribute to variation in pollutant levels. Median values were seen to decrease for all hotspots for PM<sub>2.5</sub> and PM<sub>10</sub> but increased for Jahangirpuri in the case of NO<sub>2</sub>. This could be attributed to the presence of several inter-state goods carriers in the vicinity due to a major national highway located nearby.

### EFFECT OF LOCKDOWN IN MAJOR NCR TOWNS

The air pollution reduction trend in NCR towns was relatively less pronounced compared to NCT of Delhi. Over 48% Reduction in PM<sub>10</sub> and PM<sub>2.5</sub> levels were observed during lockdown period in all neighboring towns with sharp improvement in Faridabad with 55% reduction in PM<sub>2.5</sub> levels and Gurugram with 54% reduction in PM<sub>10</sub> levels While significant reduction in NO<sub>2</sub> levels was observed in Noida (68%), Ghaziabad (60%) and Gurugram (40%), the same was not noted in Faridabad (17%), where NO<sub>2</sub> emissions were found higher during a few days in lockdown period, seemingly due to the gas-based power plants in and around Faridabad. Significant reduction in SO<sub>2</sub> levels was only seen in Faridabad (47%) and Ghaziabad (22%), while Gurugram (14%) and Noida (10%) recorded slight reductions during the lockdown period which may be attributed to their proximity to thermal power plants and some operational industries in the vicinity. Moreover, while Delhi's industries have largely switched over to gas-based and other less polluting energy systems, some industries in NCR might still be using coal and biomass, etc. Overall, average CO levels decreased in all major NCR towns with peak hourly values decreasing by 56% in Faridabad, 52% in Noida and 48% in Gurugram indicating reduced vehicular exhaust impact during lockdown. In terms of Benzene levels, Noida and Ghaziabad recorded an enormous reduction of 60% and 90% respectively during the lockdown period. However, Faridabad saw increase in Benzene levels. The operation of certain units or


processes (chemical/pharmaceutical/paints) utilizing benzene, and other solvents, etc. in Faridabad cannot be ruled out and may be responsible for the increase in Benzene levels. It is also important to mention that there are lesser number of real time air quality monitoring stations in NCR towns as compared to Delhi and the impact of localized sources on air quality data is always a possibility which may require further data for complete analysis. Moreover, in absence of complete data on scale of industrial operation in various categories except power plants and essential activities like food, bakery, dairy etc, it may be difficult to assess the impact of these sources on air quality at this point of time.




The trends for neighboring NCR towns are presented in detail in subsequent paras,




## FARIDABAD




The impact of restrictions was visible in Faridabad. 19 days in the 22-day lockdown period witnessed 24 hourly PM<sub>2.5</sub> levels within National Ambient Air Quality Standards (NAAQS). However, SO<sub>2</sub> and NO<sub>2</sub> values remained within National standards during pre-lockdown and lockdown period. While there has been a considerable drop in the peak hourly values of PM<sub>2.5</sub> (44%), PM<sub>10</sub> (60%), SO<sub>2</sub> (57%) and CO (56%) during the lockdown period as compared to the pre-lockdown period, the minimum hourly value of NO<sub>2</sub> dropped only by 10% and hourly NO<sub>2</sub> values roughly remained the same throughout the day, in all probability due to localized combustion activities and operation of gas-based power plants in the vicinity. However, 57% reduction in peak hourly SO<sub>2</sub> levels with the minimum hourly SO<sub>2</sub> levels also decreasing by 29% during the lockdown period indicates that SO<sub>2</sub> emissions from coal-based power plants in NCR might be playing a more dominant role in Delhi than Faridabad


The data trends for Faridabad is as presented below,















## GURUGRAM


PM<sub>2.5</sub>, NO<sub>2</sub> and SO<sub>2</sub> levels remained below National Ambient Air Quality Standards on all days during the lockdown, while PM<sub>10</sub> levels were above NAAQS on just 4 days in the 22-day lockdown period. Hourly data reveals a declining trend in pollutant levels from 07:00 Hrs onwards for major pollutants. Further, since dust & construction activities contribute 49% to PM<sub>2.5</sub> and 52% to PM<sub>10</sub> in Gurugram (TERI Source Apportionment study, 2018), it is likely that road dust resuspension due to vehicle restrictions might have come down resulting in lower emissions with 42% and 52% reduction in peak hourly PM<sub>2.5</sub> and PM<sub>10</sub> levels respectively. Peak hourly CO values reduced by almost 48% while peak hourly NO<sub>2</sub> and benzene levels reduced by 42% and 36% respectively during the lockdown period, indicating reduced vehicle movement. Peak SO<sub>2</sub> values fell by 20%, i.e. on a similar scale as Delhi.

The data trend for Gurugram is as presented below,







### 24 Hourly Average Benzene Comparison






### Hourly PM<sub>2.5</sub> Comparison




### Hourly PM<sub>10</sub> Comparison




















## GHAZIABAD

SO<sub>2</sub> and NO<sub>2</sub> levels remained below National Ambient Air Quality Standards on all days during the lockdown, while PM<sub>2.5</sub> levels were above NAAQS in the second week of April i.e. for 7 days in the 22-day lockdown period. Although PM<sub>2.5</sub> and PM<sub>10</sub> levels were higher during early morning hours and late-night hours, characteristically due to reduced ventilation and mixing height, peak hourly PM<sub>2.5</sub> and PM<sub>10</sub> levels reduced by 49% and 57% respectively. Major reduction of 91% in peak benzene levels and 66% in peak NO<sub>2</sub> values was observed, largely due to the reduced presence of vehicular and industrial activity. Major reduction in benzene levels (highest in Delhi NCR) during lockdown period indicate closure of some large-scale benzene utilizing/generating source like paint, petro products, plastics, resins, synthetic fibers, rubber lubricants, dyes, detergents, drugs and pesticides in Ghaziabad region apart from reduced impact of vehicular related emissions. While average CO values during lockdown period remained below their pre-lockdown levels to a great extent and reduced by almost 30%, analysis of hourly CO values indicate peak hourly CO value rising by 11%, seemingly due to local combustion activities which may include increased use of solid fuels/biomass in household cooking etc. Further, the diurnal cycle of CO concentration presents two peaks, in the morning and in the evening. Notably, mixing height is generally low in these two periods of the day. Further, studies indicate that in the early hours of the morning, surface heating by solar radiation is also not enough to break the previous night's thermal inversion layer, causing the pollutants to remain concentrated in regions close to the surface. As mixing height increases allowing transport of pollutants to the upper layers, CO levels on the surface decrease and rise again by the end of the afternoon, when convective activity decreases and traffic generally increases.














## NOIDA




Positive effects of lockdown on air pollution levels were observed in Noida, as emission levels considerably reduced from the pre-lockdown period with over 48% reduction in PM<sub>2.5</sub> and PM<sub>10</sub>. 24 hourly average PM<sub>2.5</sub> and PM<sub>10</sub> concentrations remained within NAAQS for 19 and 17 days respectively out of the 22 days in the lockdown period. NO<sub>2</sub> and SO<sub>2</sub> levels remained within NAAQS on all days of the lockdown period with the peak hourly NO<sub>2</sub> value decreasing from 76 µg/m<sup>3</sup> in the pre-lockdown period to 20 µg/m<sup>3</sup> in the lockdown period. Peak hourly Benzene levels reduced by 67%, in all possibility due to the restrictions on vehicular activity and industrial operations. While construction activity is a major emission source in Noida contributing 47% to PM<sub>10</sub> (TERI source apportionment study, 2018), significant reduction in PM<sub>10</sub> levels with hourly peak values decreasing by 55%, suggest reduced contribution of road dust resuspension & C & D activities. Reduction in PM<sub>2.5</sub> and CO emission levels was lower during morning hours signifying contribution of combustion activities. Further, over 52% reduction in peak hourly CO values was observed.




While overall SO<sub>2</sub> levels were seen to decline during the lockdown period, peak hourly SO<sub>2</sub> value increased marginally. Hourly SO<sub>2</sub> levels were also higher during the early morning hours, when dispersion of pollutants is lower. It may be said that thermal power plants located in NCR and use of fuels like coal and biomass/wood etc in industrial and household activities including operation of some brick kilns, Sugar and distilleries, might be playing a more dominant role in affecting SO<sub>2</sub> levels in Noida.











## EFFECT OF LOCKDOWN IN DELHI NCR

Substantial improvement in air quality of Delhi NCR is noted during the lockdown period, as the major contributing sources to PM & NO<sub>2</sub> emissions (prominent pollutants in Delhi NCR) have been restricted. The AQI in Delhi NCR was largely under 'moderate' category in the week before start of lockdown period. As days progressed, under cumulated effect of restricted vehicle movement, industrial & commercial activities and increased mixing height, the AQI improved to 'Satisfactory' category. On March 26, 2020, high surface winds (25 kmph) maintained AQI category even though mixing height dropped to 1100 m. Next day, though wind speed and mixing height were reduced to half value, AQI value improved further and Gurugram recorded 'Good' AQI category. Scattered rains in Delhi NCR on 27<sup>th</sup> March and during March 28- 29, 2020 along with increased wind speed and mixing height, AQI value improved further, with Delhi, Ghaziabad and Noida recording 'Good' AQI category on March 28, 2020. Favorable conditions ensued, leading to AQI remaining in 'Good' and 'Satisfactory' AQI categories. However, after 4<sup>th</sup> April due to change in temperature and onset of dry conditions, high winds led to lifting of local dust resulting in slight deterioration of air quality to 'moderate' category. Further, a dust storm from the gulf hit Delhi and the surrounding areas on 15<sup>th</sup> April, further pushing the air quality to the higher end of moderate category.



Date	Predominant Wind Speed (kmph)	Maximum Mixing Height (m)	Delhi	Ghaziabad	Noida	Faridabad	Gurugram
16-Mar	16	1800	139	134	118	184	165
17-Mar	12	1500	157	148	140	164	141
18-Mar	15	2000	151	172	137	164	168
19-Mar	14	3200	186	236	184	194	192
20-Mar	12	2400	192	235	195	212	175
21-Mar	16	2500	186	207	161	174	126
22-Mar (Janata Curfew)	12	2900	191	237	176	214	191
23-Mar	10	800	124	159	123	130	91
24-Mar	10	2700	122	166	130	187	127
<b>IMPOSITION OF NATIONWIDE LOCKDOWN DUE TO COVID-19</b>							
25-Mar	12	2500	77	86	80	100	69
26-Mar	25	1100	92	84	72	88	61
27-Mar	15	500	69	72	60	75	42
28-Mar	14	2250	45	39	38	64	54
29-Mar	20	2600	62	48	58	83	62
30-Mar	20	2100	71	64	61	97	76
31-Mar	12	1900	76	72	67	110	77
01-Apr	12	3200	73	79	73	90	69
02-Apr	20	3050	69	63	62	63	72
03-Apr	22	2100	79	104	72	97	82




<b>04-Apr</b>	20	3000	87	109	70	90	89
<b>05-Apr</b>	10	3500	102	124	84	117	91
<b>06-Apr</b>	15	3500	142	181	120	123	106
<b>07-Apr</b>	32	3750	90	101	78	100	86
<b>08-Apr</b>	20	3100	83	113	85	103	91
<b>09-Apr</b>	16	2400	86	86	80	103	96
<b>10-Apr</b>	15	3500	118	115	93	117	104
<b>11-Apr</b>	8	3050	124	194	146	203	152
<b>12-Apr</b>	20	2750	94	93	87	119	98
<b>13-Apr</b>	18	3700	126	132	120	122	106
<b>14-Apr</b>	12	4480	130	145	123	146	113
<b>15-Apr</b>	12	4980	155	194	184	186	142

## AIR QUALITY TREND ANALYSIS IN OTHER CITIES DURING LOCKDOWN PERIOD



### MUMBAI

During the lockdown period, significant reduction in PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub> levels were observed. Overall, 40% reduction in PM<sub>2.5</sub> and 43% reduction in PM<sub>10</sub> was observed during the lockdown period. A sharp decrease of 77% reduction in NO<sub>2</sub> levels and 59% decrease in average Benzene levels was observed during the lockdown period, compared to the week before lockdown came into force, largely due to the reduced presence of vehicular activity and restricted industrial operations. Average SO<sub>2</sub> levels decreased by 33% in the lockdown period, largely due to shutdown of industries. As per emission inventory of Mumbai (CPCB, 2010), 39 types of industries (excluding power plants) contribute to over 50% of SO<sub>2</sub>. 24 hourly average PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub> and NO<sub>2</sub> levels were within National Ambient Air Quality Standards for all days in the lockdown period. Lowest 24 hourly average PM<sub>10</sub>, PM<sub>2.5</sub> and NO<sub>2</sub> levels during the lockdown period were recorded as 54 µg/m<sup>3</sup>, 20 µg/m<sup>3</sup> and 5 µg/m<sup>3</sup> respectively.







The graphs below depict hourly concentration trend for PM<sub>2.5</sub> and PM<sub>10</sub>, for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).



The hourly comparison of average concentration values shows a declining trend in levels of PM<sub>10</sub> and PM<sub>2.5</sub> during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM<sub>10</sub> was 175 µg/m<sup>3</sup> at 08:00 Hrs, which dropped to 85 µg/m<sup>3</sup> during the lockdown period. Similarly, the lowest concentration during the pre-lockdown period at 17:00 Hrs was 98 µg/m<sup>3</sup>, which dropped to 58 µg/m<sup>3</sup> during the lockdown period, seemingly due to restriction on construction activities, less road dust resuspension and to some extent curb on industrial activities. A similar decline was seen for PM<sub>2.5</sub> with concentration value falling from a peak of 62 µg/m<sup>3</sup> at 08:00 Hrs (during the pre-lockdown period) to a minimum value of 21 µg/m<sup>3</sup> at 05:00 Hrs during the lockdown period. The absence of non-essential vehicles and combustion activities in industrial and commercial sites during the period may be attributable to the decline. Notably, the

lowest values are occurring in early morning hours during the lockdown period against a deeper crest in evening hours during the pre-lockdown period. This indicates the absence of nighttime accumulation of pollutants, highlighting the absence of major emission sources.



The graphs below depict hourly concentration trend for NO<sub>2</sub> and CO for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).



Hourly NO<sub>2</sub> and CO values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO<sub>2</sub> during the pre-lockdown period was almost five times the peak value observed during the lockdown period. Similar trend was observed for CO with 61% reduction in peak hourly values during the lockdown period, underscoring the absence of vehicular emissions.

The graphs below depict hourly concentration trend for SO<sub>2</sub> and Benzene for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).










As a result of the lockdown, hourly Benzene and SO<sub>2</sub> levels remained below their pre-lockdown levels at all times. The diurnal variation in the pollutants was reduced, implying the reduced impact of major emission sources. Peak hourly SO<sub>2</sub> values declined by 46% while peak hourly Benzene Concentration reduced by over 70%, seemingly due to absence of vehicular activity.


## KOLKATA


During the lockdown period, over 40% reduction in PM<sub>2.5</sub>, 44% in PM<sub>10</sub> levels and 38% reduction in SO<sub>2</sub> levels were observed. CO average levels also decreased in a similar manner with 39% reduction during the lockdown period. However, 66% reduction in NO<sub>2</sub> levels and 69% reduction in Benzene levels was observed during the lockdown period, indicating the reduction in number of on-road vehicles. 24 hourly average PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub> and NO<sub>2</sub> levels were within National Ambient Air Quality Standards for all days in the lockdown period. Lowest PM<sub>10</sub>, PM<sub>2.5</sub> and NO<sub>2</sub> 24 hourly average levels during the lockdown period were recorded as 38 µg/m<sup>3</sup>, 18 µg/m<sup>3</sup> and 6 µg/m<sup>3</sup>.






The graphs below depict hourly concentration trend for PM<sub>2.5</sub> and PM<sub>10</sub>, for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).






The hourly comparison of average concentration values shows a declining trend in levels of PM<sub>10</sub> and PM<sub>2.5</sub> during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM<sub>10</sub> was 150 µg/m<sup>3</sup> at 00:00 Hrs, which dropped to 73 µg/m<sup>3</sup> during the lockdown period. Similarly, the lowest concentration during the pre-lockdown period at 16:00 Hrs was 92 µg/m<sup>3</sup>, which dropped to 56 µg/m<sup>3</sup> during the lockdown period. The drop in coarse particles may be attributed to restriction on construction activities, less road dust resuspension and to some extent curb on industrial activities. A similar decline was seen for PM<sub>2.5</sub> with concentration value falling from a peak of 80 µg/m<sup>3</sup> at 00:00 Hrs (during the pre-lockdown period) to a minimum value of 33 µg/m<sup>3</sup> at 11:00 Hrs during the lockdown period which may be attributed to the absence of non-essential vehicles and combustion activities in industrial and commercial sites. Kolkata's diurnal variation of pollutants is characterized by lower concentrations during the day and accumulation of pollutants during nighttime which was reduced due to the absence of daytime activities like vehicular movement and industrial operation.


The graphs below depict hourly concentration trend for NO<sub>2</sub> and CO for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).






Hourly NO<sub>2</sub> and CO values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO<sub>2</sub> during the pre-lockdown period was more than thrice the peak value observed during the lockdown period. Similar trend was observed for CO, with hourly peak CO values reducing by almost 59%.

The graphs below depict hourly concentration trend for SO<sub>2</sub> and Benzene for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).










Due to restrictions on industrial operations and vehicular activity, SO<sub>2</sub> and Benzene levels remained below their pre-lockdown levels during the lockdown period. A decrease of 49% was observed in peak hourly SO<sub>2</sub> values. There was little diurnal variation in Benzene levels with over 77% reduction in peak hourly Benzene values while hourly SO<sub>2</sub> levels increased as the day progressed, which may be attributed to the thermal power plants running in and around Kolkata along with local combustion activities.

## BENGALURU



Significant reduction in PM<sub>10</sub> and NO<sub>2</sub> levels were observed during the lockdown period, as a result of combination of reduced vehicles on the road and functioning of only essential commercial units. Overall, 44% reduction in PM<sub>10</sub> and 21% reduction in PM<sub>2.5</sub> was observed during the lockdown period. Higher reduction in PM<sub>10</sub> levels indicate closure of dust generating activities like construction and Demolition etc. Due to the restrictions imposed on vehicular movement and industrial activity, a 53% reduction in NO<sub>2</sub> levels and 48% reduction in Benzene levels was observed during the lockdown period, compared to the week before lockdown came into force. 24 hourly average PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub> and NO<sub>2</sub> levels were within National Ambient Air Quality Standards for all days in the lockdown period. Lowest 24 hourly average PM<sub>10</sub>, PM<sub>2.5</sub> and NO<sub>2</sub> levels during the lockdown period were recorded as 20 µg/m<sup>3</sup>, 15 µg/m<sup>3</sup> and 9 µg/m<sup>3</sup>. However, only 15% reduction was seen in SO<sub>2</sub> levels during the lockdown period. It is possible that use of fuels like coal and biomass/wood etc in industrial and household activities, might be influencing SO<sub>2</sub> levels in Bengaluru. Similar reduction was seen in CO levels too, seemingly due to local combustion activities.







The graphs below depict hourly concentration trend for PM<sub>2.5</sub> and PM<sub>10</sub>, for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).







The hourly comparison of average concentration values shows a declining trend in levels of PM<sub>10</sub> and PM<sub>2.5</sub> during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM<sub>10</sub> was 135 µg/m<sup>3</sup> at 20:00 Hrs, which dropped to 56 µg/m<sup>3</sup> during the lockdown period. Similarly, the lowest hourly concentration during the pre-lockdown period at 03:00 Hrs was 64 µg/m<sup>3</sup>, which dropped to 37 µg/m<sup>3</sup> during the lockdown period, in all likelihood due to restriction on industrial operations, construction activities and reduced road dust resuspension. A decline was also seen for PM<sub>2.5</sub> with concentration value falling from a peak of 46 µg/m<sup>3</sup> at 20:00 Hrs (during the pre-lockdown period) to a minimum value of 18 µg/m<sup>3</sup> at 14:00 Hrs during the lockdown period, possibly due to the reduced presence of vehicles and industrial/commercial combustion activities.

The graphs below depict hourly concentration trend for NO<sub>2</sub> and CO for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).



Hourly NO<sub>2</sub> and CO values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO<sub>2</sub> during the pre-lockdown period was thrice the peak value observed during the lockdown period. Diurnal variation in NO<sub>2</sub> has reduced due to decreased vehicular activity. Hourly CO values also remained below their pre-lockdown values for most of the hours except during early morning, with hourly peak value decreasing by 30%. Post lockdown, higher CO values during night hours have reduced and maximum concentration of 1.4 mg/m<sup>3</sup> was observed at 20.00 hours before lockdown and 0.99 mg/m<sup>3</sup> at 09.00 hours during lockdown.

The graphs below depict hourly concentration trend for SO<sub>2</sub> and Benzene for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).








Peak hourly Benzene levels recorded a decrease of 47% in the lockdown period, largely due to the reduction in the number of on-road vehicles. However, there was only a 9% reduction in peak SO<sub>2</sub> values. Hourly SO<sub>2</sub> values during the lockdown period appear to follow a similar pattern as pre-lockdown levels, implying that the major sources of SO<sub>2</sub> in Bengaluru might still be operational, despite the lockdown and requires further investigation.

**PATNA**



Overall, 27% reduction in PM<sub>2.5</sub>, 27% reduction in PM<sub>10</sub> and 30% reduction in NO<sub>2</sub> levels was observed during the lockdown period, compared to the week before lockdown came into force. SO<sub>2</sub> and NO<sub>2</sub> levels remained within National Ambient Air Quality Standards for all of the days in the lockdown period. While PM<sub>2.5</sub> levels were over NAAQS on just three days, PM<sub>10</sub> levels remained within NAAQS for only 5 days in the 22-day lockdown period. Since major source of PM<sub>10</sub> for Patna is dust, it is difficult to assess its

contribution during the period without additional data like meteorology. Lowest PM<sub>10</sub>, PM<sub>2.5</sub> and NO<sub>2</sub> 24 hourly average levels during the lockdown period were recorded as 84 µg/m<sup>3</sup>, 28 µg/m<sup>3</sup> and 17 µg/m<sup>3</sup>. Benzene levels dropped by almost 61% in the lockdown period, indicating the absence of major industrial activities utilizing/emitting benzene. Although lower pollutant concentrations were observed in Patna, major reduction was not seen probably due to continuing movement of vehicles. CO and SO<sub>2</sub> levels, however were observed to slightly increase (5%), seemingly due to increased local combustion activities and usage of solid fuels in household and industrial units. However more information may be required about Sulphur dioxide sources in and around the city to examine the reasons behind the trend of SO<sub>2</sub> during lockdown.







The graphs below depict hourly concentration trend for PM<sub>2.5</sub> and PM<sub>10</sub>, for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).





The hourly comparison of average concentration values shows a declining trend in levels of PM<sub>10</sub> and PM<sub>2.5</sub> during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM<sub>10</sub> was 220 µg/m<sup>3</sup> at 22:00 Hrs, which dropped to 159 µg/m<sup>3</sup> during the lockdown period. Similarly, the lowest concentration during the pre-lockdown period at 15:00 Hrs was 117 µg/m<sup>3</sup>, which slightly reduced to 92 µg/m<sup>3</sup> during the lockdown period. A similar decline was seen for PM<sub>2.5</sub> with concentration value falling from a peak of 84 µg/m<sup>3</sup> at 23:00 Hrs (during the pre-lockdown period) to a minimum value of 32 µg/m<sup>3</sup> at 17:00 Hrs during the lockdown period, probably due to the absence of non-essential vehicles and industrial operations during the period.

The graphs below depict hourly concentration trend for NO<sub>2</sub> and CO for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).



Hourly NO<sub>2</sub> and CO values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO<sub>2</sub> during the pre-lockdown period was over one and half times the peak value observed during the lockdown period. Though 11% reduction in hourly peak CO values was observed in the lockdown period, overall levels of CO were seen to increase, probably due to local combustion activities like biomass burning and agri-waste burning in surrounding areas.

The graphs below depict hourly concentration trend for SO<sub>2</sub> and Benzene for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).





While peak hourly Benzene value was seen to decrease by 61% in the lockdown period, peak hourly SO<sub>2</sub> value increased by 3%. Hourly SO<sub>2</sub> values during the lockdown period appear to be following the same diurnal variation as in the pre-lockdown period, implying that the major sources of SO<sub>2</sub>, i.e. biomass burning or usage of unclean fuels in household and industrial activities including brick kilns might still be operational. The trend of SO<sub>2</sub> need to be further investigated with more information on likely sources including on-ground data, as it is defying general trend observed during lockdown.



## INDORE

The impact of lockdown in Indore was prominent especially for PM<sub>10</sub>, NO<sub>2</sub> and CO which saw 29%, 56% and 40% reduction during the lockdown period respectively, seemingly due to the reduction in the number of on-road vehicles. However, the reduction in PM<sub>2.5</sub> was only about 5%, compared to the week before lockdown came into force. It is important to mention that there is only one real time air quality monitoring





stations in Indore and the impact of localized sources on air quality data is always a possibility. Gradual increase in PM<sub>2.5</sub> and PM<sub>10</sub> in the second week of April may be attributed to dust and operation of some combustion sources using coal/biomass. SO<sub>2</sub> and NO<sub>2</sub> levels remained within National Ambient Air Quality Standards for all of the days in the lockdown period. While PM<sub>2.5</sub> levels were over NAAQS on just four days, PM<sub>10</sub> levels remained within NAAQS for 14 days in the 22-day lockdown period. Lowest PM<sub>10</sub>, PM<sub>2.5</sub> and NO<sub>2</sub> 24 hourly average levels during the lockdown period were recorded as 27 µg/m<sup>3</sup>, 12 µg/m<sup>3</sup> and 22 µg/m<sup>3</sup>. Benzene levels were not available in the CAAQM station. SO<sub>2</sub> levels also decreased by 34%, which may be attributed to the restriction on industrial operations.







The graphs below depict hourly concentration trend for PM<sub>2.5</sub> and PM<sub>10</sub>, for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).




The hourly comparison of average concentration values shows a declining trend in levels of PM<sub>10</sub> during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM<sub>10</sub> was 261 µg/m<sup>3</sup> at 21:00 Hrs, which dropped to 141 µg/m<sup>3</sup> during the lockdown period. Similarly, the lowest concentration during the pre-lockdown period was 89 µg/m<sup>3</sup> at 14:00 Hrs, which reduced to 67 µg/m<sup>3</sup> during the lockdown period. A decline was seen for PM<sub>2.5</sub> with concentration value falling from a peak of 91 µg/m<sup>3</sup> at 21:00 Hrs (during the pre-lockdown period) to a minimum value of 18 µg/m<sup>3</sup> at 13:00 Hrs during the lockdown period, probably due to the absence of non-essential vehicles and restriction on industrial operations during the period. However, the hourly values were slightly higher during the early morning hours, and requires further data for further analysis.

The graphs below depict hourly concentration trend for NO<sub>2</sub> and CO for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).



Hourly NO<sub>2</sub> values during the lockdown period remained below the hourly values observed during the pre-lockdown period. The peak hourly value of NO<sub>2</sub> during the pre-lockdown period was over two and a half times the peak value observed during the lockdown period, indicating the reduced presence of vehicular activity. Though 66% reduction in hourly peak CO values was observed in the lockdown period, CO levels were higher in the early morning hours, probably due to local combustion activities. Further, mixing height is less during early morning, thereby inhibiting dispersion of pollutants.




The graph below depict hourly concentration trend for SO<sub>2</sub> for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).




While peak hourly SO<sub>2</sub> value was seen to decrease by 66% in the lockdown period, SO<sub>2</sub> values during the lockdown period were higher for a few hours after midnight when compared to the levels in the pre-lockdown period. This may be due to reduced dispersion at night-time and needs further data.

## CHENNAI

Chennai saw over 36% reduction in PM<sub>2.5</sub>, seemingly due to restriction on vehicular activities and reduced combustion activities. The reduction in NO<sub>2</sub> and CO levels was around 5% and 43% during the lockdown period. Studies have reported that while transport sector is the prominent source for CO, industries contribute majorly to NO<sub>x</sub> levels. Since two of the CAAQM stations are located in Manali, Chennai which is an industrial area having significant presence of chemical and petrochemical industries, it is possible that the industrial operations in the area might be influencing the NO<sub>2</sub> levels in Chennai. Further, this is corroborated by the fact that vehicles and industries are the two major sources of NO<sub>x</sub> (CPCB, 2010) in Chennai. PM<sub>2.5</sub>, SO<sub>2</sub> and NO<sub>2</sub> levels remained within National Ambient Air Quality Standards for all of the days in the lockdown period and PM<sub>10</sub> data was not available for Chennai. Lowest PM<sub>2.5</sub> and NO<sub>2</sub> 24 hourly average levels during the lockdown period were recorded as 12 µg/m<sup>3</sup> and 4 µg/m<sup>3</sup> while SO<sub>2</sub> levels decreased by 29%.





The graph below depicts hourly concentration trend for PM<sub>2.5</sub>, for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).



The hourly comparison of average concentration values shows a declining trend in levels of PM<sub>2.5</sub> during the lockdown period. During the pre-lockdown period, the maximum hourly value of PM<sub>2.5</sub> was 84 µg/m<sup>3</sup> at 14:00 Hrs, which dropped to 34 µg/m<sup>3</sup> during the lockdown period. However, the sudden spike in PM<sub>2.5</sub> in the pre-lockdown period does seem unnatural. Similarly, the lowest concentration during the pre-lockdown period was 26 µg/m<sup>3</sup> at 02:00 Hrs, which reduced to 19 µg/m<sup>3</sup> during the lockdown period, probably due to the absence of non-essential vehicles and restriction on industrial operations during the period. However, the hourly values were slightly higher for one hour after midnight, analysis of which requires further data.


The graphs below depict hourly concentration trend for NO<sub>2</sub> and CO for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).



Hourly NO<sub>2</sub> values during the lockdown period kept fluctuating and were both above and below the hourly values observed during the pre-lockdown period at different hours of the day. The peak hourly value of NO<sub>2</sub> during the lockdown period was just 6% less than the peak value observed during the pre-lockdown period, while hourly peak CO levels reduced by 46%. This indicates that while vehicular and local combustion activity may have reduced, some industries might still be operational in the area. Further, there is little diurnal variation, implying the absence of major sources of CO.

The graph below depict hourly concentration trend for SO<sub>2</sub> for pre-lockdown period (16<sup>th</sup> March 2020 to 21<sup>st</sup> March 2020) and lockdown period (25<sup>th</sup> March 2020 to 15<sup>th</sup> April 2020).





Peak hourly SO<sub>2</sub> values were seen to decrease by 18% in the lockdown period, seemingly due to restriction on industrial operations. Hourly SO<sub>2</sub> values during lockdown period were less than the concentration observed during pre-lockdown period for most hours, with the only deviations occurring after midnight, which may be due to adverse meteorology.

#### **EFFECT OF LOCKDOWN IN OTHER CITIES IN TERMS OF AIR QUALITY INDICES**

Air Quality Indices are calculated for cities all over India using data from CAAQM stations. If cities appearing in CPCB AQI Bulletin are grouped according to their respective AQI categories, it is observed that about 78% of cities in the AQI bulletin are falling in Good and Satisfactory categories in the lockdown phase, increasing from the average of 44% seen in the pre-lockdown phase. Since Good and Satisfactory categories have their breakpoints within the National Ambient Air Quality Standards, it may be reasonable to state that more cities have their air quality within National standards during the lockdown period.

During the lockdown period, no city entered the very poor category. Among the cities in poor category during the lockdown period, instances of Singrauli and Brajrajnagar are found frequently. It is worth noting that Singrauli region is home to several power plants, which are operational during the lockdown period and Brajrajnagar has in its vicinity numerous open-cast and underground coal mines.

The date wise AQI is given in Annexure I.

## IMPACT OF LOCKDOWN ON AIR QUALITY –SATELLITE BASED MONITORING

Satellite-based monitoring is being used widely for air quality data generation. The advantage of satellite-based estimates is that they can provide country wide coverage of all locations. A project entitled “Satellite based near real time monitoring of ambient PM<sub>2.5</sub> at national scale for air quality management” has been initiated by CPCB in collaboration with Indian Institute of Technology (IIT), Delhi under National Clean Air Programme (NCAP) in order to strengthen development of indigenous satellite-based products and techniques to derive useful air quality information and to supplement the current monitoring network.

Impact of lockdown on PM<sub>2.5</sub> levels at a national scale was carried out using Aerosol Optical Depth for the period March 18 to April 13, 2020. Weekly PM<sub>2.5</sub> concentration anomaly (in %) in 2020 has been reported relative to 2015-2019 average concentration (see Figure). The analysis suggests that PM<sub>2.5</sub> was higher in North and Central India during the pre-lockdown week this year. With many emission sources shut down, in the first two weeks of lockdown, PM<sub>2.5</sub> was 30-50% lower in most parts of India, especially in the Indo-Gangetic Plain (IGP).

However, this improvement was not as prominent in the 2<sup>nd</sup> week of April, when the dust influence started to dominate. Data after 13<sup>th</sup> April is being analysed to understand whether this is temporary trend, however, ground-based data has showed that air quality is again back in moderate zone.




Figure. Anomaly (in % w.r.t last 5-years) in weekly surface PM<sub>2.5</sub> from satellite-derived data during pre- and lockdown period

## **SUMMARY**

During the lockdown period, there has been a general improvement in air quality in the country as a result of the restrictions imposed during the lockdown, which is corroborated by both Air Quality Index data and Satellite data. All of the major cities, analysed in this report, had their AQI, largely within Good-Moderate categories, with Bengaluru, Chennai and Mumbai recording all 22 days of the lockdown period in Good-Satisfactory categories while Patna had majority of days in moderate category.

**Comparative AQI Status from 16 March to 15 April, 2020  
(based on CPCB AQI Bulletin, published at 4 PM)**

Date	No of cities for which data is available	No. of cities in AQI category						No. of cities with AQI in range of Good to Satisfactory	No. Of cities with AQI in Moderate Category	No. of cities with AQI in range of Poor to Severe
		Good	Satisfactory	Moderate	Poor	Very Poor	Severe			
16-Mar-20	108	6	49	50	3	0	0	55	50	3
17-Mar-20	111	3	44	59	5	0	0	47	59	5
18-Mar-20	112	3	42	58	9	0	0	45	58	9
19-Mar-20	115	3	39	65	8	0	0	42	65	8
20-Mar-20	115	2	51	50	12	0	0	53	50	12
21-Mar-20	112	2	52	49	9	0	0	54	49	9
22-Mar-20 (Janata Curfew)	114	9	58	39	8	0	0	67	39	8
23-Mar-20	108	10	63	33	2	0	0	73	33	2
24-Mar-20	110	11	54	43	2	0	0	65	43	2
<b>National Lockdown in effect due to COVID-19 Pandemic</b>										
25-Mar-20	104	14	67	21	2	0	0	81	21	2
26-Mar-20	102	21	64	14	3	0	0	85	14	3
27-Mar-20	103	31	59	10	3	0	0	90	10	3
28-Mar-20	101	35	57	8	1	0	0	92	8	1
29-Mar-20	103	30	61	12	0	0	0	91	12	0
30-Mar-20	99	23	65	11	0	0	0	88	11	0
31-Mar-20	103	20	67	14	2	0	0	87	14	2
01-Apr-20	100	23	62	13	2	0	0	85	13	2
02-Apr-20	105	22	71	11	1	0	0	93	11	1

<b>03-Apr-20</b>	105	20	71	14	0	0	0	<b>91</b>	<b>14</b>	0
<b>04-Apr-20</b>	109	22	68	18	1	0	0	<b>90</b>	<b>18</b>	1
<b>05-Apr-20</b>	104	17	65	21	1	0	0	<b>82</b>	<b>21</b>	1
<b>06-Apr-20</b>	102	23	49	29	1	0	0	<b>72</b>	<b>29</b>	1
<b>07-Apr-20</b>	101	25	56	18	2	0	0	<b>81</b>	<b>18</b>	2
<b>08-Apr-20</b>	102	22	54	25	1	0	0	<b>76</b>	<b>25</b>	1
<b>09-Apr-20</b>	102	20	58	24	0	0	0	<b>78</b>	<b>24</b>	0
<b>10-Apr-20</b>	104	17	54	31	2	0	0	<b>71</b>	<b>31</b>	2
<b>11-Apr-20</b>	103	21	49	29	4	0	0	<b>70</b>	<b>29</b>	4
<b>12-Apr-20</b>	108	17	62	28	1	0	0	<b>79</b>	<b>28</b>	1
<b>13-Apr-20</b>	104	15	50	38	1	0	0	<b>65</b>	<b>38</b>	1
<b>14-Apr-20</b>	102	8	53	36	5	0	0	<b>61</b>	<b>36</b>	5
<b>15-Apr-20</b>	105	8	49	38	10	0	0	<b>57</b>	<b>38</b>	10

<b>AQI Category</b>	<b>AQI Range</b>	<b>Associated Health Impact</b>
<b>Good</b>	0-50	Minimal Impact
<b>Satisfactory</b>	51-100	Minor breathing discomfort to sensitive people
<b>Moderate</b>	101-200	Breathing discomfort to the people with lungs, asthma and heart diseases
<b>Poor</b>	201-300	Breathing discomfort to most people on prolonged exposure
<b>Very Poor</b>	301-400	Respiratory illness on prolonged exposure
<b>Severe</b>	401-500	Affects healthy people and seriously impacts those with existing diseases

**Number of cities with AQI in Good, Satisfactory and Moderate categories**

